

(2)

EUROPEAN PATENT APPLICATION

(31) Application number: 89300231.1

(51) Int. Cl.⁴: **A61K 47/00 , A61K 39/012**

(22) Date of filing: 11.01.89

Claims for the following Contracting States: ES
+ GR.

(30) Priority: 11.01.88 US 141953

(43) Date of publication of application:
26.07.89 Bulletin 89/30

(54) Designated Contracting States:
AT BE CH DE ES FR GB GR IT LI LU NL SE

(71) Applicant: **A.H. ROBINS COMPANY,
INCORPORATED**
1407 Cummings Drive P.O. Box 26609
Richmond Virginia 23261-6609(US)

(72) Inventor: **Bhogal, Balbir Singh**
12091 Cottage Mill Place
Midlothian Virginia 23113(US)

(74) Representative: **Sheard, Andrew Gregory et al**
Kilburn & Strode 30, John Street
London WC1N 2DD(GB)

(54) **Live vaccine for coccidiosis utilizing coccidial sporozoites.**

(57) Methods and compositions are disclosed for vaccinating warm-blooded animals against coccidiosis utilizing suspensions of excysted coccidial sporozoites in physiologically balanced medium containing water-soluble polymeric stabilizers selected from gels, gelatins, polysaccharide gums, cellulose or cellulose derivatives which extend viability or storage, additional extension of viability in storage being attained when the suspensions are finely divided and the polymeric stabilizers are hardened to form microcapsules.

EP 0 325 359 A1

LIVE VACCINE FOR COCCIDIOSIS UTILIZING COCCIDAL SPOROZOITES

This invention relates to methods of vaccinating warm-blooded animals against coccidiosis disease and compositions therefor.

Biology and Life Cycle of Coccidial Protozoa

1. Coccidial protozoa may infect and inhabit the gastrointestinal tract of most species of warm-blooded animals, most often and most seriously when the animals are penned or in concentrated groups. For the most part infection occurs with coccidial species specific to the animal host species and more than one coccidial species may infect a given host animal species. For example, seven major coccidial species are known to infect the chicken.

The life cycle of the coccidial parasite involves several complex stages. Under natural conditions the cycle begins with oral ingestion by the host animal of excrement from another animal of the same species having sporulated oocysts dispersed therein. The sporulated oocyst shell then breaks down by action of the digestive system of the host animal. For example when the host animal is a chicken the outer shell of the sporulated oocyst is broken down mechanically to liberate first the sporocysts and finally biochemically to liberate sporozoites which are the ultimate infective organism. Other stages in the cycle which follow redevelopment within about 7 days include schizont I, merozoite I, schizont II, merozoite II, gametocyte, gamete and completion of the cycle with oocyst formation.

More descriptively, in the instance of the chicken, natural infection with one of the important species of *Eimeria*, i.e., *Eimeria tenella* begins with ingestion of an overwhelming amount of the sporulated oocysts. Each sporulated *E. tenella* oocyst contains four sporocysts and each sporocyst contains two sporozoites. In the stomach, the sporulated oocysts rupture, freeing the sporocysts. The sporozoites are liberated from the sporocysts by enzymatic action and pass down the digestive tract. Each of the sporozoites eventually reach the cecal pouches where they enter epithelial cells of the cecal mucosa and start an asexual multiplication cycle. The details of the completion of the life cycle referred to above are well known in the art.

The rationale of vaccination with live sporozoites in the present invention, is to set up a low level of infection (subclinical) during the first few days of life of the animal, e.g., a chicken, without slowing down the performance of the animal in terms of growth. The subclinical infection, thus established induces immunity, while the host, e.g., chicken, has the advantages of passive protection afforded by maternal antibodies. The maternal antibodies also help to down regulate infection, thus helping to decrease unnecessary parasite buildup in animal houses, which is critical for the success of a vaccine since the degree of severity of disease is directly proportional to the numbers of sporulated oocysts ingested by the host from its environment. Under field conditions chickens, e.g., have their maternal antibodies reduced to very low levels by the third week and hence become susceptible to the parasite, if their immune system is not primed by vaccination. Without vaccination the challenge encountered by hosts is frequently devastating, especially to concentrated populations of animals resulting in death or illness and/or poor development or growth quality and subsequent economic loss to the animal grower.

U.K. Patent Specification No. 2,008,404 describes a method of promoting the growth of poultry by feeding viable sporulated oocysts of coccidia to poultry. Similarly chickens have been said to be immunized against cecal coccidiosis with viable sporulated oocysts in the disclosure of U.S. Patent 3,147,186.

An immunogenic composition for oral administration comprising live encysted protozoa such as coccidia is described in British Application 2,144,331A wherein encysted sporulated coccidial oocysts or encysted coccidial sporocysts embedded in a firm gel matrix were fed orally to the host animal such as a newly-hatched poultry chick. Gels based on alginate were used for the matrix.

Sharma, N.N. in *The Journal of Parasitology* Vol. 50, No. 4 (1984) pp. 509-517 describes unsuccessful immunization trials with sporulated viable oocysts of coccidia or with an impure mixture said to contain sporozoites along with oocysts and sporocysts of coccidia administered to chickens, orally, intravenously, intraperitoneally, intramuscularly or subcutaneously.

Since the severity of disease is directly proportional to the number of sporulated oocysts in the field available for ingestion, existing vaccines based on either feeding sporulated oocyst or feeding microencapsulated oocyst have limited utility for the broiler chicken industry, in that the high numbers of oocysts available for ingestion from day one often causes severe infections. As a result, the use of oral vaccines as such often necessitates the use of anticoccidials. Therefore the use of sporulated oocysts as live vaccines

have limited application. Additionally, sporozoites are ineffective in producing an infection when administered orally to chickens.

Heretofore unsuccessful attempts to create immunity with unprotected sporozoites by systemic inoculation have been made by Sharma, supra., where attempts to create infection with sporozoites have been confined to use of freshly excysted sporozoites inasmuch as methods have not previously been available to keep the sporozoites alive longer than a few days under refrigeration. Consequently vaccination using sporozoites was impractical commercially.

Therefore according to a first aspect of the present invention there is provided a composition for vaccinating warm-blooded animals, such as fowl and in particular chickens, against coccidiosis disease comprising an optionally microencapsulated suspension of live excysted coccidial sporozoites in aqueous medium, physiologically balanced with respect to isotonicity and pH, and containing at least one water-soluble viscosity enhancing polymeric stabiliser. Such compositions may serve to prolong the viability of the sporozoites.

The sporozoites are usually *Eimeria* sporozoites, such as those of the species *Eimeria tenella*.

The polymeric stabiliser is preferably a hydrosol, gel, gelatin, cellulose, cellulose derivative, polysaccharide gum or alginate. Sodium alginate is particularly preferred and in the case of microencapsulation, the sporozoites are suitably microencapsulated by calcium alginate.

According to a second aspect of the invention there is provided a vaccine comprising microencapsulated excysted coccidial sporozoites. As in the first aspect, the sporozoites can be *Eimeria* sporozoites, such those of the species *Eimeria tenella*.

The sporozoites may be encapsulated by polymeric stabiliser such as hydrosol, gel, gelatin, cellulose, cellulose derivative, polysaccharide gum or alginate but particularly preferred is calcium alginate.

A third aspect of the present invention relates to a veterinary composition comprising a composition or vaccine in accordance with the first or second aspects and a verterinarily acceptable carrier.

A fourth aspect of the present invention concerns the use of optionally microencapsulated coccidial sporozoites and at least one water-soluble viscosity enhancing polymeric stabilizer in the preparation of a vaccinating agent. The use extends to vaccination of fowl, in particular chicken, subcutaneously such as in the back of the neck.

A fifth aspect of the present invention relates to a process for the preparation of a composition or veterinary composition of the first and third aspects, such as a composition for vaccinating warm-blooded animals against coccidiosis disease, comprising admixing a suspension of live excysted coccidial sporozoites, water and at least one water-soluble viscosity enhancing polymeric stabilizer, wherein the sporozoites are optionally microencapsulated.

A sixth aspect relates to a process for the preparation of a vaccine in accordance with the second aspect, the process comprising microencapsulating the sporozoites.

A preferred method of this invention of microencapsulating by finely dividing a suspension and hardening the water-soluble polymeric stabilizer to enclose a physiologically balanced suspension of sporozoites may provide time flexibility as to when the vaccine must be used, since the lifetime (in terms of viability and infectivity) of the sporozoites may be extended up to at least 5 weeks. A second method of this invention of using water-soluble polymeric stabilizer, preferably a water-soluble alginate salt to suspend sporozoites in a medium physiologically balanced with respect to isotonicity and pH may extend the lifetime of the sporozoites for up to at least 2 weeks and thus also may provide time flexibility as to when the vaccine must be used.

The invention encompasses microencapsulated excysted live coccidial sporozoites suspended in stabilizing carriers which may greatly extend life expectancy or shelf-life of the sporozoites. More particularly aqueous compositions containing water soluble polymeric stabilizers, and physiologically balanced as to isotonicity and pH for all components which are used as suspending medium and carriers for the sporozoites. The water soluble polymeric stabilizers used may be classed as gels, gelatins, cellulose, cellulose derivatives, and polysaccharide gums, especially water soluble alginate salts. Preferably these sporozoite suspensions containing a water soluble polymeric stabilizer are first finely divided and then the surface of the particles is hardened to form microcapsules leaving an internal suspension of sporozoites in the stabilizing aqueous medium.

The technique of microencapsulation of sporozoites used here may not only provides a practical way of vaccination of chickens when they are handled on their first day of life, but also offers other advantages, such as, e.g., avoiding build up of oocysts in the animal yard or pen which occurs both during vaccination as well as afterwards due to parasite multiplication in the host when live oocysts are used commercially, i.e., spread in the yard or pen or used in feed to inoculate, and thus avoids early overwhelming challenge due to that source of oocysts. Use of alginate suspension and microcapsules containing suspensions of

sporozoites, in contrast, results in much slower and lower levels of oocyst buildup underfoot.

The invention also encompasses a suspension of excysted coccidia sporozoites, physiologically balanced with respect to isotonicity and pH, containing sodium alginate in solution, and having a viscosity in the range of from 100 to 400 centipoises at 25°.

5 The sporozoites of Eimeria species once out of their protective shells, i.e., oocysts and sporocysts, are very fragile and lose their infectivity within a few days. This invention is based on the discovery that Eimeria sporozoites can be kept alive for at least 5 weeks by microencapsulation and that the longevity of live excysted sporozoites can be increased to at least 2 weeks by addition of buffered aqueous suspensions of the sporozoites to solutions of water soluble polymeric stabilizers including sodium alginate. The microen-
 10 capsulation of live sporozoites provides a buffered viscous aqueous medium for the parasite consisting of a solution of water soluble polymeric stabilizer, said stabilizer preferably being sodium alginate. The microcapsules containing the aged sporozoites can be administered to baby chicks to establish subclinical infections thereby vaccinating against any future overwhelming challenge of E. tenella oocysts. Similar effective vaccination of chickens substituting other species of coccidia natural to the chicken and similar
 15 vaccination of other animals including poultry substituting viable excysted coccidial sporozoites appropriate to the animal species are reasonable expectations. Similar effective vaccinations with unencapsulated sporozoite buffered suspensions containing the water soluble polymeric stabilizers are also viable compositions.

Purified, excysted live coccidial sporozoites are first obtained in a suspension of isotonic buffered
 20 aqueous solution by known procedures which involve mechanical, chemical and enzymatic breakdown of sporulated coccidial oocysts and sporocysts to remove the sporecase covers therefrom followed by chromatographic purification. Isotonic buffered sporozoite suspensions are added to solutions of protective agents (isotonic buffered polymeric stabilizers) which delay the death of the sporozoites and thereby extend the length of time the sporozoites remain viable in storage, such suitable protective agents being selected
 25 from hydrocolloids, gelatins, cellulose, cellulose derivatives and polysaccharides, especially alkali-metal alginate salts. Preferably the suspension so obtained is finely divided as a spray and mixed with hardening agent to give microcapsules which surround or contain the suspension of live excysted sporozoites in isotonic buffered medium.

According to the vaccination method of the invention, the isotonic buffered sporozoite suspension
 30 containing the protective agent, microencapsulated or unencapsulated, is administered to warm blooded animals to protect against severe damage to intestinal organs due to overwhelming challenge of coccidial oocysts. The microencapsulated preparations may be stored for at least 5 weeks. Although the preferred method is microencapsulation, the unencapsulated preparations may be stored for at least 2 weeks before using when kept refrigerated but not frozen. Young animals, are treated with preparations carrying a
 35 quantified number of the excysted microencapsulated sporozoites appropriately chosen for the animal species involved. While any route of administration may be used, the subcutaneous route such as in the back of the neck, is preferred particularly for fowl, for example chickens. Vaccines and compositions of the present invention to be administered to warm blooded animals preferably sterile.

Certain terms used in the present discussion should at this point be mentioned to avoid ambiguity of
 40 their respective meanings. Thus the following definitions are provided to clarify the scope of the present invention and to enable its formulation and use.

An "oocyst" is the dormant life-cycle stage of the coccidial protozoa having a tough outer coat. In its initial formation the core of the oocyst is in an immature state not having as yet elements of infective capability and may be referred to as an "unsporulated oocyst." Such "oocysts" are found in the gut of
 45 animals prior to elimination or on the ground just after elimination.

A "sporulated oocyst" is an oocyst which has undergone biochemical maturation brought about naturally or artificially under certain temperature and oxygen exposure conditions. One common artificial sporulating condition is exposure to dichromate solution. During maturation a multiplicity of "sporocysts" each within its own case or shell develops within the outer case thus becoming "cysts within a cyst." In the
 50 instance of the chicken, e.g., E. tenella, have four separate encased sporocysts within the outer shell after the oocyst have undergone sporulation. Live sporulated oocysts are commonly referred to as "viable sporulated oocysts."

"Encysted coccidial protozoa," "encysted oocysts" and "encysted sporocysts" all refer to organisms which are within the cyst or have their own natural shell or sporecase.

55 Each sporocyst shell surrounds the ultimate infective agent, the "sporozoite." In the instance of E. tenella each sporocyst shell encompasses two sporozoites.

The term "excysted sporozoite" refers to the live sporozoite from which the protecting shell of the sporocyst has been removed. Excystation occurs naturally by biochemical reaction but is accomplished, for

use of the excysted sporozoites, in the method of this invention by homogenization of a suspension of pure sporulated oocysts, centrifuging to remove solubles, treating the solids with a solution of trypsin and taurodeoxycholic acid in Hank's solution at pH 7.4 followed by purification using anion exchange chromatography as described more fully infra. Methods of obtaining sporozoites, as described herein
 5 bypass isolation of sporocysts as such.

"Capsule Wall Material" refers to outer hardened surface of the microcapsule which results from cross-linking of the water soluble polymeric stabilizer with a suitable cross linking agent. Such material preferably calcium alginate.

Suitable "cross-linking agents" are divalent cations in the instance of polysaccharide gums, and
 10 aldehydes such as glutaraldehyde or formaldehyde in the instance of gels and gelatins.

"Water soluble polymeric stabilizers" or "Polymeric stabilizers" and "viscosity enhancing substances" all refer generally to gels, gelatins, cellulose, cellulose derivatives and polysaccharide gums, one or more of which are required in the compositions of this invention. Included in this group of materials are alginates of high and low viscosity, acacia, agar, agarose, albumen, dextran, various gelatins, polyvinyl alcohol, sodium
 15 carboxycellulose and starches.

The specific way in which these materials function to stabilize the sporozoites and increase longevity is not known, however, a suitable viscosity range for prolonging sporozoite stability (viability) in suspensions of this invention is 100-400 centipoises as measured at about 25° C.

The term "modified capsule wall" as used herein refers to the microcapsule shell, e.g., calcium
 20 alginate, having been formed which has been treated after formation, usually after storage and prior to administration to allow easier escape of sporozoites from the microcapsule. Suitable capsule wall modifiers for the alginate system are salts of weak acids such as citric acid, fumaric acid and maleic acid.

By the use of the term "physiologically balanced," when used in conjunction with the compositions herein described is meant the compositions are physiologically balanced with respect to isotonicity and pH
 25 of living cells, the pH being in the range of 6.8 to 7.5; preferably about 7.2 and having osmolality in the range of 260-350 milliosmoles. The suspension of sporozoites are thus buffered to physiological pH and osmotically balanced with respect to living cells. Generally amine buffers may be used to physiologically balance, for example, TRIS buffer and HEPES buffer are effective buffering agents.

The term "suspension" as used herein refers to both liquid and solid suspensions, should the latter
 30 occur within the microcapsule. Microcapsules (or microspheres) usually have a diameter of from 300 to 400 microns, more usually from 340 to 360 microns.

Compositions useful in the vaccination procedure of the invention may contain sporozoites of a single strain or species of coccidia; however when broad host-specific immunity is required, a selection of sporozoites of common strains or species of coccidia should be included in a single formulation.

The invention may be applied to the control of coccidiosis in any animal host species that is prone to
 35 the disease eg. poultry. In addition to the avian species, pigs and ruminants such as cattle, sheep, goats and rabbits are affected by the disease and may benefit from use of this method of this invention.

In chickens about seven major species of coccidial parasitize the intestinal tract beginning with the duodenum; they are: Eimeria acervulina, E. mivati, E. Maxima, E. necatrix, E. brunetti, E. mitis, and E. tenella. In turkeys, the following coccidia have been observed: E. melagrimitis, E. disperse, E. meleagridis,
 40 E. gallopavonis, E. adenoides, E. innocus and E. subrotunda. In pigs the following coccidia have been observed: E. debliciecki, E. scabra, E. suis, E. spinosa, E. perminuta, E. neodebliciecki, E. porci, E. cordonis, E. polita and Isospora suis. In cattle the following coccidia have been observed: E. bovis, E. zuernii, E. ellipsoidalis, E. auburnensis, E. cylindrica, E. alabamensis and E. bukidnonensis. In sheep, the following
 45 coccidia have been observed: E. minokohlyakimovae, E. ovina, E. intricate and E. aheate. Goats are infected by E. arloinge and rabbits are infected by E. intestinalis, E. flavesens, E. magna, E. irrisidua, E. periformis, E. stiedai, E. performans, E. neoleporis and E. media.

50 2 Preparation 1. Excysted E. tenella Sporozoites In Buffer Solution

2.1 General Procedure for Isolating Excysted Sporozoites

The procedure used to obtain purified Eimeria tenella sporozoites was that of Schmatz, D.M. *et al.* described in J. Protozol 31(1)(1984) pp. 181-183 hereby and described stepwise in an abbreviated form
 55 following. In addition microscopic observations was employed from time to time.

2.2 Obtaining Precursor Sporulated Oocysts

Coalesced masses of oocysts were isolated by scraping cecal pouches of chickens which had been infected seven days earlier with *Eimeria tenella* (A. H. Robins strain 855). The cecal cores in distilled water were disrupted in a Waring Blender® and digested with pepsin at pH 2.0 and 39 °C for 1 hr. Debris and pepsin were removed by centrifugation in distilled water. A partially purified oocyst fraction was isolated by flotation in 1.1 molar aqueous sucrose solution. This semi-pure oocyst fraction was incubated in cold Chlorox® solution for 15 min at 4 °C. The Chlorox® component was removed by washing several times in sterile phosphate buffered saline (PBS) at pH 7.3. The resulting purified sterile viable oocysts were suspended in dichromate solution and then sporulated by shaking in a water bath at 29 °C for 48 hr. The sporulated oocysts in PBS were stored at 4 °C until used in the next step.

2.3 Excystation of Sporozoites and Buffered Solution Thereof

A 2 ml suspension of PBS (pH 7.3) as prepared above containing sporulated oocysts, was ground in a tissue homogenizer operated at about 500 rpm for 5 min at 4 °C. Soluble material resulting from disruption of the sporulated oocysts was removed by centrifugation. The pellet obtained, composed of unbroken oocysts, sporocysts and oocyst shells, was resuspended in excysting solution which contained as excysting agents: 0.25% (weight/volume) trypsin (1:250) and 4% (weight/volume) taurodeoxycholic acid (Sigma Co., St. Louis, Missouri) in Hank's balanced salt solution (pH 7.4). The mixture was incubated in an atmosphere of 5% carbon dioxide at 41 °C for 1 hr. The excysting solution was removed by two washes with stock column buffer solution. Column buffer solution for this and subsequent column purification was prepared as follows: A stock buffer solution of 2 x phosphate-buffered saline (2 x PBS, 26.96 g/liter disodium phosphate, 1.56 g/liter disodium phosphate, 8.5 g/liter sodium chloride) was used to prepare a solution 2:8 (vol/vol) PBS:H₂O of 0.145 ionic (I) strength.

2.4 Chromatographic Purification of Excysted Sporozoites

The excysted sporozoites in buffer solution were purified using anion exchange column chromatography in specially prepared cellulose (DE-52 cellulose) packed column as described by Schmatz, *ibid*, the appearance of the solution under the microscope confirming the completeness of the purification.

DE-52 cellulose anion exchange column packing was prepared by equilibrating pre-swollen Whatman (England) DE-52 cellulose in proportions of 2.0 g in 150 ml of the 2:8 buffer (I=0.145). After settling, the supernatant containing the fines was removed. This washing procedure was repeated twice more and the DE-52 cellulose was adjusted to pH=8.0 with 5% (wt/vol) phosphoric acid prior to the last wash. This material was then resuspended in 50 ml of the 2:8 buffer and used to fill the column.

The columns used to purify excysted sporozoites were sterilized by autoclaving prior to filling. Columns were packed to a height of 2.5 cm with the DE-52 cellulose, as prepared above, and the packing was rinsed with 50 ml of 2:8 buffer (I=0.145).

Freshly excysted sporozoites as prepared above in 2:8 buffer were loaded onto a column loaded with 2:8 equilibrated DE-52 cellulose, as prepared above. Portions of 2:8 buffer were sequentially passed through the column. The portions were collected off the column and the content evaluated and quantified by hemocytometer under phase microscopy. The suspension of excysted sporozoites in buffered solution thus obtained was suitable for adding to solutions containing stabilizers described herein such as sodium alginate solutions.

3 Preparation 2. Suspension of Excysted *E. tenella* Sporozoites in Isotonic Buffered Aqueous Sodium Alginate Solution

The following is a stepwise description of preparation of alginate-stabilized suspensions of sporozoites suitable as vaccine to be used directly or microencapsulated.

3.1 TRIS Buffer Solution

0.1 M solution of TRIS Buffer [Tris (hydroxymethyl)aminomethane] was prepared by accurately weighing 6.057 g of TRIS crystals and diluting with 500 ml of distilled water.

5 3.2 Buffered Sodium Alginate Solution

1.5% sodium alginate (low viscosity) solution was prepared by dissolving 7.5 g of sodium alginate in 500 ml of TRIS buffer prepared above. The solution is cooled to 4 °C.

10

3.3 Calf Serum Addition

Freshly prepared concentrated suspension having 1×10^8 sporozoites in 5 ml of TRIS buffer was obtained (from Preparation 1) and 5 vol % fetal calf serum was added and the suspension was kept at 4 °C.

15

3.4 Buffered Sporozoite-Alginate Suspension

The 5 ml of buffered solution containing the 1×10^8 sporozoites in 5.3.3 above was added to 145 ml of the alginate solution prepared in 5.3.2 above with stirring until blending was complete. The suspension of sporozoites was then maintained at 4 °C. These proportions were based on microencapsulation needs in Preparation 3 wherein calculations are explained.

25 4 Preparation 3. Microcapsules Containing Stabilized Excysted E. tenella Sporozoites in Isotonic Buffered Aqueous Solution

The following is a description of the procedure for encapsulating the stabilized suspension of excysted sporozoites:

30

4.1 Calculations on Volume Ratios

The proportion of buffered sodium alginate solution to buffered sporozoite suspension used to prepare the product of Preparation 2 was based on calculation to produce microcapsules of a certain size using the formula below using the following facts and assumptions: 1) that 5 ml of buffered sporozoite suspension contained 1×10^8 sporozoites, 2) that microcapsules of 350 micron diameter would be prepared, 3) that it was the aim to have each microcapsule contain 15 sporozoites and 4) that based on the radius of 175 microns and the formula for calculating the volume of a sphere ($\frac{4}{3} \pi r^3$) there would be 44,545 microcapsules per ml of final alginate solution.

40

$$1 \times 10^8 \text{ sporozoites} \times \frac{1 \text{ microcap}}{15 \text{ sporozoites}} \times \frac{1 \text{ ml alginate suspension}}{44,545 \text{ microcapsules}} = 150 \text{ ml of final alginate suspension}$$

45

Therefore 145 ml of the alginate solution was added to 5 ml of buffered solution containing the 1×10^8 sporozoites prepared in Preparation 2.

50

4.2 Preparation of Hardening Bath

A hardening bath of 1.5% calcium chloride solution was prepared by dissolving 19.63 g of calcium chloride dihydrate (76.4% calcium chloride) in 1 liter of distilled water. The osmolality of this solution was about 335 milliosmoles/kg.

55

4.3 Description of Gas Flow Droplet Generator (Jet-Spray Head)

The jet spray head used to finely divide the sodium alginate sporozoite suspension was mounted above the hardening bath and perpendicularly lengthwise to the bath surface.

The jet-spray head consisted of a housing having a small elongated conical chamber 3/4" in height x 1 1/4" ID. at the top and 1/16" ID at the bottom, capped with a rubber disc at the top broader end and open at the narrower end, the chamber having a side gas port and a liquid feed needle protruding through the cap and downward through the length of the chamber so that the open end of the needle extended well into the chamber. A pump was used to regulate liquid flow through the needle and a gas pressure valve was used to control gas flow into the chamber via the sideport. The needle used was a 21 gauge x 1 1/2" syringe needle with a conically shaped tip.

10

4.4 Preparation of Microcapsules Containing Sporozoites Suspended in Stabilizing Aqueous Solution

With the bottom end of the jet-spray head set about 1.5 cm above the hardening bath, the sporozoite-alginate suspension from Preparation 2 under constant agitation at 4°C was pumped with a peristaltic pump at a flow rate of 0.46 ml/min against a gas flow through the side port of about 8 liters per min, the microspheres being blown off into the hardening bath. The size of the microspheres was adjusted to about 350 micron diameter, using microscopic examination of the microcapsules being obtained to determine the size, by manipulating the gas flow rate and the position of the needle opening. The microspheres immediately hardened on entering the bath. This batch of microcapsules utilized 135 ml of the starting 150 ml of sporozoite suspension described above to give a total of 1×10^6 microcapsules.

20

4.5 Washing of Microcapsules with Saline

25

The microcapsules prepared above were allowed to settle to a stable volume in the hardening bath and excess calcium chloride was decanted off. The residual microcapsules, wet with calcium chloride solution were resuspended in an equivalent volume of normal saline, i.e., aqueous 0.9 wt % sodium chloride solution, at 4°C and allowed to settle again. The washing process was repeated three times. The microcapsules in normal saline were then refrigerated at about 4°C until they were used as vaccine or modified and used as vaccine as in paragraph 5.9 below.

30

4.6 Storage Tests Relative to Viability of Microencapsulated Sporozoites

35

The microcapsules as prepared in 5.4.5 in normal saline were stored for 5 weeks at 4°C. The saline was decanted and the microcapsules were resuspended in PBS solution (pH = 7.3). The suspension was administered subcutaneously in the back of the neck to one week old Hubbard x Hubbard male chicks. The chicks were sacrificed 7 and 8 days after administration and cecal pouches found to be infected with coccidia. It is concluded therefore that sporozoites contained in a suspension of physiologically compatible medium containing an alginate salt, the whole of which suspension of sporozoites is microencapsulated by crosslinking the alginate with a divalent cation, remained viable for at least 5 weeks when stored in normal saline at 4°C.

40

45

5.5 Example 1. Demonstration of Sub-Clinical Infection With Microencapsulated and Unencapsulated Sporozoites of E. tenella In Chickens

A short initial study affirmed that low-level infection could be obtained with both unencapsulated live excysted sporozoites as obtained in Preparation 2, and microencapsulated live excysted sporozoites as obtained in Preparation 3. Four groups of one-day old chickens were sacrificed after 7 or 8 days exposure and their cecal pouches were examined for lesions. The study is summarized in Table I.

50

55

Table I
Demonstration of Infectability With Microencapsulated Sporozoites

Group No.	Number of Chickens in Group	Sporozoite Treatment		Cecal Infection Observed on Day:	
		Kind	Method of Administration(a)	Day 7	Day 8
1	15	microcaps, 10 ⁵ live excysted sporozoites	S.Q. (back of neck)	lesions 7/8	lesions 8/8
2	9	10 ⁵ live excysted sporozoites, no encapsulation	S.Q. (back of neck)	lesions 7/9	.
3	9	10 ⁵ live excysted sporozoites, no encapsulation	i.p.	lesions 8/9	.
4	10	none (controls)		no lesions 10/10	.

S.Q. = subcutaneously on back of neck

i.p. = intraperitoneally

Example 2. Demonstration of Vaccination Potential Against E. tenella Coccidia Using Microencapsulated Excysted Stabilized Sporozoites In Chickens

Two hundred and forty (240) one-day old Hubbard x Hubbard broiler chicks were divided into four groups, chickens being randomly selected so that their body weights were within 10% of the mean. Groups 1 and 2 were in replicates of 2 floor pens each pen housing 40 chicks for a total of 80 chicks in each of groups 1 and 2. Groups 3 and 4 were in single pens housing 40 chicks per pen. Each group of chicks were individually injected subcutaneously (S.Q.) in the back of the neck on day one as follows:

Group 1:

Each chick received S.Q. microencapsulated PBS-sodium alginate suspension of 10^5 live excysted sporozoites, the microcapsules being suspended in normal saline, volume totalling 0.1 ml. The microcapsules were those prepared in Preparation 3.

Group 2:

Each chick received S.Q., microencapsulated PBS-sodium alginate suspension of 5×10^3 live excysted sporozoites, the microcapsules being suspended in normal saline, volume totalling 0.1 ml. The microcapsules were those prepared in Preparation 3.

Group 3:

Each chick received subcutaneously, PBS suspension of 10^5 live freshly excysted sporozoites in 0.1 ml volume. See Preparation 1 for procedure of obtaining sporozoites. No microencapsulation was involved.

Group 4:

Each chick received 0.1 ml of PBS solution only.

Twenty eight days later each chick was fed an oral challenge of 85,000 E. tenella oocysts.

On Day 34 (i.e. 6 days after the oocyst challenge) the chicks were sacrificed and both cecal pouches of each chick were examined for damage to the cecal walls using the scoring procedure of Joyce Johnson and W. Malcolm Reid described in Experimental Parasitology 28, 30-36 (1970) on page 31 for E. tenella with grading scores applicable as follows:

Grading Score:

- 0 No gross lesions
- +1 Very few scattered petechiae on the cecal wall; no thickening of the cecal walls; normal cecal contents present.
- +2 Lesions more numerous with noticable blood in the cecal contents; cecal wall is somewhat thickened; normal cecal contents present.
- +3 Large amounts of blood or cecal cores present; cecal walls greatly thickened; little if any fecal contents in the ceca.
- +4 Cecal walls greatly distended with blood or large caseous cores; fecal debris lacking or included in cores. Dead birds scored as +4.

Table II summarizes important features of the test and shows net cumulative results of the test on each group as a percentage of all individuals exhibiting a low scoring range 0-1.5, a middle scoring range 2-2.5 or a high scoring range 3-4. As will be readily realized by one skilled in the art, a high percentage in the low scoring range of 0-1.5 is desirable and is indicative of the effectiveness of the vaccination procedure.

Table II

Immunisation Studies With Microencapsulated Excysted <i>E. tenella</i> Sporozoites In Chickens (Example 2 Summary)				
Group No.	No. of Chickens Per Group	Day 1: Sporozoite Treatment Subcutaneous	Day 28: Oocyst Challenge Oral	Day 34: Cecal Damage 8 days after Oocyst Challenge, % of Chickens in Scoring Ranges of:
				0-1.5 (low) 1.75-2.5 3-4 (high)
1	2 pens of 40 each	microencapsulated PBS alginate suspension of 10^5 sporozoites	85,000 Sporulated oocysts	42 18 40
2	2 pens of 40 each	microencapsulated PBS alginate suspension of 5×10^8 sporozoites	85,000 Sporulated oocysts	11 10 79
3	1 pen of 40	freshly prepared suspension of 10^5 sporozoites	85,000 Sporulated oocysts	93 - 5
4	1 pen of 40	PBS 0.1 ml vol (no sporozoites)	85,000 Sporulated oocysts	9 16 75

Subcutaneous in back of neck

Microcapsules in 0.1 ml vol. with normal saline

Example 3. Demonstration of Vaccination Potential Against E. tenella Coccidia Using Microencapsulated Excysted Stabilized Sporozoites In Chickens

5 Three hundred and sixty (360), one-day old Hubbard x Hubbard broiler chicks were divided into 18 pens of 20 chicks each, randomly selected so that their body weights were within .10% of the mean. The eighteen groups were randomly divided into 3 pens each for 6 treatment groups. Groups 1-5 were adjacently located and Group 5 was in a separate area. The treatment groups were as follows. Treatment
10 for each group on day one was as follows:

Group 1:

15 Each chick in 3 pens received, S.Q., 2 week old microencapsulated PBS suspension of 1×10^5 live excysted E. tenella sporozoites, the microcapsules being suspended in normal saline, volume totalling 0.1 ml. The microcapsules were prepared by a procedure similar to that of Preparation 3, of about 250 micron average size.

20 Group 2:

Same as Group 1 except administration was intraperitoneally (i.p.) to chicks in 3 pens.

25 Group 3:

Freshly prepared 1×10^5 excysted E. tenella sporozoites - PBS suspension was administered S.Q., to chicks in 3 pens (0.1 ml).

30 Group 4:

Same as group 3, except suspension of sporozoites was administered intraperitoneally to chicks in 3
35 pens (0.1 ml).

Group 5:

40 0.1 ml of PBS solution only was administered S.Q., to chicks in 3 pens.

Group 6:

45 Three pens of chicks were kept isolated, no treatment given.

Twenty seven days later each chick was fed an oral challenge of 50,000 E. tenella oocysts.

On day 33 (= 6 days after the oocyst challenge) the chicks were sacrificed and both cecal pouches of each chick were examined for damage to the cecal walls the scoring procedure of Joyce Johnson and W. Malcom Reid described in Example 2 herein.

50 Table III summarizes important feature of the test and shows net cumulative results of the test on each group as a percentage of all individuals exhibiting scoring ranges of 0-1.5, 2-2.5 and 2.75-4.

Table III

Immunization Studies With Microencapsulated Excysted <i>E. tenella</i> Sporozoites In Chickens (Example 3 Summary)						
Group No.	No. of Chickens Per Group	Day 1: Sporozoite Treatment (Route)	Day 28: Oocyst Challenge Oral	Day 34: Cecal Damage 6 days after Oocyst Challenge, % of Chickens in Scoring Ranges of:		
				0-1.5	2.0-2.5	2.75-4
1	3 pens of 20 each	microencapsulated PBS suspension of 10 ⁵ sporozoites (S.Q.)	50,000 oocysts	43	38	19
2	3 pens of 20 each	microencapsulated PBS suspension of 10 ⁵ sporozoites (i.p.)	50,000 oocysts	50	25	25
3	3 pens of 20 each	10 ⁵ freshly prepared sporozoites in PBS (S.Q.)	50,000 oocysts	90	10	0
4	3 pens of 20 each	10 ⁵ freshly prepared sporozoites in PBS (i.p.)	50,000 oocysts	94	0	6
5	3 pens of 20 each	PBS Solution only (S.Q.)	50,000 oocysts	30	35	35
6	3 pens of 20 each (isolated)	None	None	5	20	75

Example 4. Stabilizing Effect of Sodium Alginate on E. tenella Sporozoites

5 To 200 ml of a 1.5% solution of sodium alginate (low viscosity) was added 1 ml of a freshly prepared suspension having 10^6 - 10^7 sporozoites/ml in TRIS buffered solution containing calf serum as prepared in par. 5.3.3 so as to give a suspension of 5,000 excysted sporozoites/ml. After 14 days storage at 4° C, chicks were injected per cloacal with 0.2 ml (1000 sporozoites) of the suspension. Infective ability was still present at that time (examination of cecal pouches for lesions) whereas sporozoites in TRIS buffer
10 containing calf serum had no infective ability after 14 days storage at 4° C.

Example 5. Vaccination Potential Using Microcapsules Having Capsule Wall Modified with Citrate

15 Microcapsules prepared from alginate as in paragraph 5.4, supra, and suspended in isotonic solution and stored at 4° C for periods of time up to 5 weeks were mixed with a 1.5% aqueous sodium citrate solution at pH 7.2 for a period of time of 1-2 minutes. Using a microscope, liberation of sporozoites was observed. The suspension, so treated, when administered subcutaneously to newly hatched chicks is expected to give improved inoculation against E. tenella oocyst challenge compared to the unmodified
20 microcapsules due to greater efficiency and speed of sporozoite release from the microcapsule.

Claims

- 25 1. A composition for vaccinating warm-blooded animals against coccidiosis disease comprising an optionally microencapsulated suspension of live excysted coccidial sporozoites in aqueous medium, physiologically balanced with respect to isotonicity and pH, and containing at least one water-soluble viscosity enhancing polymeric stabiliser.
- 30 2. A composition as claimed in Claim 1 wherein the sporozoites are of the species Eimeria tenella.
3. A composition as claimed in Claim 1 or 2 wherein the water-soluble viscosity enhancing polymeric stabiliser is a hydrosol, gel, gelatin, cellulose, alginate.
4. A composition as claimed in any of Claims 1 to 3 wherein the water-soluble viscosity enhancing polymeric stabiliser is sodium alginate and the suspension of live sporozoites are optionally microencapsulated by calcium alginate.
- 35 5. A vaccine comprising microencapsulated excysted coccidial sporozoites.
6. A vaccine as claimed in Claim 5 wherein the sporozoites are Eimeria sporozoites.
7. A vaccine as claimed in Claim 5 or 6 wherein the sporozoites are microencapsulated by a hardened water soluble polymer.
8. A vaccine as claimed in any of Claims 5 to 7 wherein the sporozoites are encapsulated by a
40 hydrosol, gel, gelatin, cellulose, cellulose derivative, polysaccharide gum or alginate.
9. A veterinary composition for vaccinating warm-blooded animals against coccidiosis disease comprising an optionally microencapsulated suspension of live excysted coccidial sporozoites in aqueous medium, physiologically balanced with respect to isotonicity and pH, and containing at least one water-soluble viscosity enhancing polymeric stabilizer; and a veterinarily acceptable carrier.
- 45 10. A composition as claimed in Claim 9 wherein the sporozoites are of the species Eimeria tenella.
11. A veterinary composition comprising microencapsulated excysted coccidial sporozoites and a veterinarily acceptable carrier.
12. A composition as claimed in Claim 11 wherein the sporozoites are Eimeria sporozoites.
13. The use of optionally microencapsulated coccidial sporozoites and at least one water-soluble
50 viscosity enhancing polymeric stabilizer in the preparation of a vaccinating agent.
14. The use of microencapsulated excysted coccidial sporozoites in the preparation of a vaccinating agent.
15. The use as claimed in Claim 14 wherein the sporozoites are Eimeria sporozoites.

55

Claims for the following Contracting States: ES, GR

1. The use of optionally microencapsulated coccidial sporozoites and containing at least one water-soluble viscosity enhancing polymeric stabilizer in the preparation of a vaccinating agent.
2. The use as claimed in Claim 1 wherein the sporozoites are of the species Eimeria tenella.
3. The use as claimed in Claims 1 or 2 wherein the water-soluble viscosity enhancing polymeric stabilizer is a hydrosol, gel, gelatin, cellulose, cellulose derivative, polysaccharide gum or alginate.
4. The use as claimed in any of claims 1 to 3 wherein the water-soluble polymeric stabilizer is sodium alginate and the suspension of live sporozoites are optionally microencapsulated by calcium alginate.
5. The use as claimed in any of claims 1 to 4 wherein the vaccinating agent is suitable for vaccinating chickens.
6. The use of a vaccine comprising microencapsulated excysted coccidial sporozoites in the preparation of a vaccinating agent.
7. The use as claimed in Claim 6 wherein the sporozoites are Eimeria sporozoites.
8. The use as claimed in Claim 6 or 7 wherein the sporozoites are microencapsulated by a hardened water soluble polymer.
9. The use as claimed in any of Claims 6 to 8 wherein the sporozoites are encapsulated by a hydrosol, gel, gelatin, cellulose, cellulose derivative, polysaccharide gum or alginate.
10. The use as claimed in any of Claims 6 to 9 wherein the vaccine is administered to a chicken.
11. The use as claimed in any of claims 6 to 10 wherein the sporozoites are of the species Eimeria tenella.
12. A process for the preparation of composition for vaccinating warm-blooded animals against coccidiosis disease comprising admixing a suspension of live excysted coccidial sporozoites, water, and at least one water-soluble viscosity enhancing polymeric stabilizer, to produce a physiologically balanced preparation with respect to isotonicity and pH, wherein the sporozoites are optionally microencapsulated.
13. A process as claimed in Claim 12 wherein the sporozoites are of the species Eimeria tenella.
14. A process for the preparation of a vaccine sporozoites, the process comprising encapsulating the sporozoites.
15. A process as claimed in Claim 14 wherein the sporozoites are Eimeria sporozoites.



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number

EP 89 30 0231

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. CL.4)
D,X	FR-A-2 408 351 (UNILEVER N.V.) * Page 6, lines 10-34; page 7, lines 19-28; page 8, lines 18-23; page 12, lines 1-7; page 14, lines 34-38; claims 1,8-15 * & GB-A-2 008 404 ---	1-3,5-15	A 61 K 47/00 A 61 K 39/012
D,X	GB-A-2 144 331 (UNILEVER PLC) * Page 1, lines 19-30,34-36,43-47; page 2, lines 32,65; page 5, lines 12-15; claims * -----	1-15	
			TECHNICAL FIELDS SEARCHED (Int. CL.4)
			A 61 K
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 24-04-1989	Examiner BERTE M.J.
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons ----- & : member of the same patent family, corresponding document			